

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/000654 A3

(51) International Patent Classification⁷: C07K 1/36.
C07D 241/08 // A61K 47/00

(21) International Application Number: PCT/US00/17984

(22) International Filing Date: 29 June 2000 (29.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/141,433 29 June 1999 (29.06.1999) US

(71) Applicant: PHARMACEUTICAL DISCOVERY CORPORATION [US/US]; 33 West Main Street, Elmsford, NY 10523 (US).

(72) Inventors: STEINER, Solomon, S.; 24 Old Wagon Road, Mount Kisco, NY 10549 (US). WOODS, Rodney, J.; 1213 Ridgebury Road, New Hampton, NY 10956 (US). SULLNER, Joseph, W.; 330 Bullard Avenue, Patamcus, NJ 07652 (US).

(74) Agents: PABST, Patrea, L. et al.; Holland & Knight LLP, One Atlantic Center, Suite 2000, 1201 West Peachtree Street, N.E., Atlanta, GA 30309-3400 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
5 July 2001

(48) Date of publication of this corrected version:
25 July 2002

(15) Information about Correction:
see PCT Gazette No. 30/2002 of 25 July 2002, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PURIFICATION AND STABILIZATION OF PEPTIDE AND PROTEINS IN PHARMACEUTICAL AGENTS

(57) Abstract: Methods are provided for purifying peptides and proteins by incorporating the peptide or protein into a diketopiperazine or competitive complexing agent to facilitate removal of one or more impurities, i.e. undesirable components, from the peptide or protein. In a preferred embodiment, a peptide, such as insulin, containing one or more impurities, e.g. zinc ions, is entrapped in diketopiperazine to form a precipitate of peptide/diketopiperazine/impurity, which is then washed with a solvent for the impurity to be removed, which is a nonsolvent for the diketopiperazine and a nonsolvent for the peptide. Formulations and methods also are provided for the improved transport of active agents across biological membranes, resulting for example in a rapid increase in blood agent concentration. The formulations include microparticles formed of (i) the active agent, which may be charged or neutral, and (ii) a transport enhancer that masks the charge of the agent and/or that forms hydrogen bonds with the target biological membrane in order to facilitate transport. In a preferred embodiment, insulin is administered via the pulmonary delivery of microparticles comprising fumaryl diketopiperazine and insulin in its biologically active form. The charge on the insulin molecule is masked by hydrogen bonding it to the diketopiperazine, thereby enabling the insulin to pass through the target membrane. This method of delivering insulin results in a rapid increase in blood insulin concentration that is comparable to the increase resulting from intravenous delivery.

WO 01/000654 A3

PURIFICATION AND STABILIZATION OF PEPTIDE
AND PROTEINS IN PHARMACEUTICAL AGENTS

Background Of The Invention

5 The present invention is generally in the field of pharmaceutical formulations, and more particularly related to methods and compositions for purifying and stabilizing peptides and proteins, such as insulin, which are used in pharmaceutical applications.

10 In a normal person, the β -cells of the pancreatic islets of Langerhans produce insulin, required by the body for glucose metabolism, in response to an increase in blood glucose concentration. The insulin metabolizes incoming glucose and temporarily stops the liver's conversion of glycogen and lipids to glucose thereby allowing the body to support metabolic activity between meals. The Type I diabetic, however, has a reduced ability or
15 absolute inability to produce insulin due to β -cell destruction and needs to replace the insulin via daily injections or an insulin pump. More common than Type I diabetes, though, is Type II diabetes, which is characterized by insulin resistance and increasingly impaired pancreatic β -cell function. Type II diabetics may still produce insulin, but they may also require insulin
20 replacement therapy.

 Type II diabetics typically exhibit a delayed response to increases in blood glucose levels. While normal persons usually release insulin within 2-3 minutes following the consumption of food, Type II diabetics may not secrete endogenous insulin for several hours after consumption. As a result,
25 endogenous glucose production continues after consumption (Pfeiffer, *Am. J. Med.*, 70:579-88 (1981)), and the patient experiences hyperglycemia due to elevated blood glucose levels.

 Loss of glucose-induced insulin secretion is one of the earliest disturbances of β -cell function (Cerasi et al., *Diabetes*, 21:224-34 (1972);
30 Polonsky et al., *N. Engl. J. Med.*, 318:1231-39 (1988)), but the causes and degree of β cell dysfunction are unknown in most cases. While genetic factors play an important role, (Leahy, *Curr. Opin. Endocrinol. Diabetes*,

2:300-06 (1995)), some insulin secretory disturbances seem to be acquired and may be at least partially reversible through optimal glucose control.

Optimal glucose control via insulin therapy after a meal can lead to a significant improvement in natural glucose-induced insulin release by requiring both normal tissue responsiveness to administered insulin and an abrupt increase in serum insulin concentrations. Therefore, the challenge presented in the treatment of early stage Type II diabetics, those who do not have excessive loss of β -cell function, is to restore the release of insulin following meals.

Most early stage Type II diabetics currently are treated with oral agents, but with little success. Subcutaneous injections of insulin are also rarely effective in providing insulin to Type II diabetics and may actually worsen insulin action because of delayed, variable, and shallow onset of action. It has been shown, however, that if insulin is administered intravenously with a meal, early stage Type II diabetics experience the shutdown of hepatic glucogenesis and exhibit increased physiological glucose control. In addition, their free fatty acids levels fall at a faster rate than without insulin therapy. While possibly effective in treating Type II diabetes, intravenous administration of insulin, is not a reasonable solution, as it is not safe or feasible for patients to intravenously administer insulin at every meal.

Insulin, a polypeptide with a nominal molecular weight of 6,000 Daltons, traditionally has been produced by processing pig and cow pancreas to isolate the natural product. More recently, however, recombinant technology has been used to produce human insulin *in vitro*. Natural and recombinant human insulin in aqueous solution is in a hexameric configuration, that is, six molecules of recombinant insulin are noncovalently associated in a hexameric complex when dissolved in water in the presence of zinc ions. Hexameric insulin is not rapidly absorbed. In order for recombinant human insulin to be absorbed into a patient's circulation, the hexameric form must first associate into dimeric and/or monomeric forms before the material can move into the blood stream. The delay in absorption

requires that the recombinant human insulin be administered approximately one half hour prior to meal time in order to produce therapeutic insulin blood level, which can be burdensome to patients who are required to accurately anticipate the times they will be eating. To overcome this delay, analogs of recombinant human insulin, such as HUMALOG™, have been developed, which rapidly disassociate into a virtually entirely monomeric form following subcutaneous administration. Clinical studies have demonstrated that HUMALOG™ is absorbed quantitatively faster than recombinant human insulin after subcutaneous administration. See, for example, U.S. Patent No. 5,547,929 to Anderson Jr., et al.

In an effort to avoid the disadvantages associated with delivery by injection and to speed absorption, administration of monomeric analogs of insulin via the pulmonary route has been developed. For example, U.S. Patent No. 5,888,477 to Gonda, et al. discloses having a patient inhale an aerosolized formulation of monomeric insulin to deposit particles of insulin on the patient's lung tissue. However, the monomeric formulation is unstable and rapidly loses activity, while the rate of uptake remains unaltered.

While it would be desirable to produce rapidly absorbable insulin derived from natural sources, transformation of the hexameric form into the monomeric form, such as by removing the zinc from the complex, yields an insulin that is unstable and has an undesirably short shelf life. It therefore would be desirable to provide monomeric forms of insulin, while maintaining its stability in the absence of zinc. It also would be advantageous to provide diabetic patients with monomeric insulin compositions that are suitable for pulmonary administration, provide rapid absorption, and which can be produced in ready-to-use formulations that have a commercially useful shelf-life.

These problems with impurities, metal ions that affect stability or bioavailability, occur with many other proteins and peptides.

U.S. Patent No. 6,071,497 to Steiner, et al. discloses microparticle drug delivery systems in which the drug is encapsulated in diketopiperazine

microparticles which are stable at a pH of 6.4 or less and unstable at pH of greater than 6.4, or which are stable at both acidic and basic pH, but which are unstable at pH between about 6.4 and 8. The patent does not describe monomeric insulin compositions that are suitable for pulmonary
5 administration, provide rapid absorption, and which can be produced in ready-to-use formulations that have a commercially useful shelf-life.

It would therefore be advantageous to develop alternative insulin delivery compositions for Type II diabetics that provide more rapid elevation of insulin blood levels and are easily administered to ensure patient
10 compliance. It also would be desirable to apply the delivery compositions and methods to other biologically active agents.

It is therefore an object of the present invention to provide improved methods for purifying peptides and proteins, especially in the preparation of compositions suitable for pulmonary administration.

15 It is another object of the present invention to provide stable monomeric peptide compositions suitable for pulmonary delivery.

It is a further object of the present invention to provide methods and compositions for the facilitated transport of insulin and other biologically active agents across biological membranes.

20 It is another object of the present invention to provide methods and compositions for the improved absorption of insulin or other biologically active agents in the bloodstream.

It is a still further object of the present invention to provide methods and compositions for the improved absorption of insulin or other biologically
25 active agents in the bloodstream characterized by ease of administration.

Summary Of The Invention

Methods are provided for purifying peptides and proteins by incorporating the peptide or protein into a diketopiperazine or competitive complexing agent to facilitate removal one or more impurities, i.e.
30 undesirable components, from the peptide or protein. In a preferred embodiment, a peptide, such as insulin, containing one or more impurities, e.g., zinc ions, is entrapped in diketopiperazine to form a precipitate of

peptide/diketopiperazine/impurity, which is then washed with a solvent for the impurity to be removed, which is a nonsolvent for the diketopiperazine and a nonsolvent for the peptide. Alternatively, the impurity can be removed by using complexing agents to selectively complex with and displace the
5 impurities, for example, such as by dialysis.

Formulations and methods also are provided for the improved transport of active agents across biological membranes, resulting, for example, in a rapid increase in blood agent concentration. The formulations include microparticles formed of (i) the active agent, which may be charged
10 or neutral, and (ii) a transport enhancer that masks the charge of the agent and/or that forms hydrogen bonds with the target biological membrane in order to facilitate transport. In a preferred embodiment, insulin is administered via pulmonary delivery of microparticles comprising fumaryl diketopiperazine and insulin in its biologically active form. The charge on
15 the insulin molecule is masked by hydrogen bonding it to the diketopiperazine, thereby enabling the insulin to pass through the target membrane. This method of delivering insulin results in a rapid increase in blood insulin concentration that is comparable to the increase resulting from intravenous delivery.

20 **Brief Description of the Drawings**

Figure 1a is a graph of mean blood glucose values over time (minutes). Figure 1b is a graph of mean C-peptide concentrations during experiments comparing levels of C-peptide (ng/ml) over time (minutes) when insulin was administered intravenously, subcutaneously, and by
25 inhalation.

Figure 2a is a graph of glucose infusion rate (mg/kg/min) over time (minutes) comparing insulin administered intravenously, subcutaneously, and by inhalation. Figure 2b is a graph of mean insulin concentrations (μ U/ml) over time (minutes) comparing insulin administered intravenously,
30 subcutaneously, and by inhalation.

Detailed Description Of The Invention

Encapsulation or entrapment of large polymers, such as proteins and peptides, in diketopiperazines can be used to remove impurities or contaminants such as metal ions or other small molecules. The
5 diketopiperazines also serve both to stabilize and enhance delivery of the entrapped materials. Formulations also have been developed for the enhanced transport of active agents across biological membranes. These formulations include microparticles formed of (i) the active agent, which may be charged or neutral, and (ii) a transport enhancer that masks the
10 charge of the agent and/or that forms hydrogen bonds with the membrane. The formulations can provide rapid increases in the concentration of active agent in the blood following administration of the formulations.

For example, it was discovered that hexameric insulin can be delivered to the lung in fumaryl diketopiperazine formulation, reaching peak
15 blood concentrations within 3-10 minutes. In contrast, insulin administered by the pulmonary route without fumaryl diketopiperazine typically takes between 25-60 minutes to reach peak blood concentrations, while hexameric insulin takes 30-90 minutes to reach peak blood level when administered by subcutaneous injection. This feat has been successfully replicated several
20 times and in several species, including humans.

Removing zinc from insulin typically produces unstable insulin with an undesirably short shelf life. Purification to remove zinc, stabilization and enhanced delivery of insulin is demonstrated by the examples. Formulations of insulin trapped in fumaryl diketopiperazine were found to be stable and
25 have an acceptable shelf life. Measurement of the zinc levels demonstrated that the zinc had been largely removed during the entrapment process, yielding monomeric insulin in a stable delivery formulation.

Rapid absorption of a number of other peptides, including salmon calcitonin, parathyroid hormone 1-34, octreotide, leuprolide and RSV
30 peptide, has been observed when the peptide is pulmonarily delivered in fumaryl diketopiperazine--providing peak blood concentrations within 3-10 minutes after pulmonary delivery.

I. Materials

A. Agent to be Delivered

The agent to be delivered is referred to herein as the active agent, or molecule to be encapsulated or entrapped. It may or may not be a charged species. Examples of classes of active agents suitable for use in the compositions and methods described herein include therapeutic, prophylactic, and diagnostic agents, as well as dietary supplements, such as vitamins.

The exact mechanism by which the diketopiperazines form a complex with the materials to be delivered is not known, but it is believed that the diketopiperazines form a complex with the material to be purified. This process is referred to herein interchangeably as entrapment or encapsulation.

These materials can be any polymer or large organic molecules, most preferably peptides and proteins. Generally speaking, any form of drug can be entrapped. Examples include synthetic inorganic and organic compounds, proteins and peptides, polysaccharides and other sugars, lipids, and nucleic acid sequences having therapeutic, prophylactic or diagnostic activities. Proteins are defined as consisting of 100 amino acid residues or more; peptide are less than 100 amino acid residues. Unless otherwise stated, the term protein refers to both proteins and peptides. The agents to be incorporated can have a variety of biological activities, such as vasoactive agents, neuroactive agents, hormones, anticoagulants, immunomodulating agents, cytotoxic agents, antibiotics, antivirals, antisense, antigens, and antibodies. In some instances, the proteins may be antibodies or antigens which otherwise would have to be administered by injection to elicit an appropriate response. Representative polymers including proteins, peptides, polysaccharides, nucleic acid molecule, and combinations thereof.

Preferred peptides and proteins include hormones, cytokines and other immunomodulatory peptides, and antigens/vaccines. In a preferred embodiment, the active agent is monomeric insulin or a stabilized form of insulin which has been purified to remove zinc. In another preferred embodiment, the active agent is glucagon.

The active agent, or drug, can be an antigen, where the molecule is intended to elicit a protective immune response, especially against an agent that preferentially infects the lungs, such as mycoplasma, bacteria causing pneumonia, and respiratory syncytial virus. In these cases, it may also be
5 useful to administer the drug in combination with an adjuvant, to increase the immune response to the antigen.

Any genes that would be useful in replacing or supplementing a desired function, or achieving a desired effect such as the inhibition of tumor growth, could be introduced using the matrices described herein. As used
10 herein, a "gene" is an isolated nucleic acid molecule of greater than thirty nucleotides, preferably one hundred nucleotides or more, in length. Examples of genes which replace or supplement function include the genes encoding missing enzymes such as adenosine deaminase (ADA) which has been used in clinical trials to treat ADA deficiency and cofactors such as
15 insulin and coagulation factor VIII. Genes which effect regulation can also be administered, alone or in combination with a gene supplementing or replacing a specific function. For example, a gene encoding a protein which suppresses expression of a particular protein-encoding gene, or *vice versa*, which induces expression of a protein-encoding gene, can be administered in
20 the matrix. Examples of genes which are useful in stimulation of the immune response include viral antigens and tumor antigens, as well as cytokines (tumor necrosis factor) and inducers of cytokines (endotoxin), and various pharmacological agents.

Other nucleic acid sequences that can be utilized include antisense
25 molecules which bind to complementary DNA to inhibit transcription, ribozyme molecules, and external guide sequences used to target cleavage by RNAase P.

As used herein, vectors are agents that transport the gene into targeted cells and include a promoter yielding expression of the gene in the
30 cells into which it is delivered. Promoters can be general promoters, yielding expression in a variety of mammalian cells, or cell specific, or even nuclear versus cytoplasmic specific. These are known to those skilled in the art and

can be constructed using standard molecular biology protocols. Vectors increasing penetration, such as lipids, liposomes, lipid conjugate forming molecules, surfactants, and other membrane permeability enhancing agents are commercially available and can be delivered with the nucleic acid.

5 Imaging agents including metals, radioactive isotopes, radioopaque agents, fluorescent dyes, and radiolucent agents also can be incorporated. Examples of radioisotopes and radioopaque agents include gallium, technetium, indium, strontium, iodine, barium, and phosphorus.

10 Impurities which can be removed from the active agent composition include metal ions such as zinc, and other di- or multi-valent ions, and small inorganic molecules and solvent residuals.

B. Diketopiperazines


Diketopiperazines useful in the present compositions and methods are described, for example, in U.S. Patent No. 6,071,497, which is incorporated
15 herein in its entirety.

(i). General Formula

The diketopiperazines or their substitution analogs are rigid planar rings with at least six ring atoms containing heteroatoms and unbonded electron pairs. One or both of the nitrogens can be replaced with oxygen to create the substitution analogs diketomorpholine and diketodioxane,
20 respectively. Although it is possible to replace a nitrogen with a sulfur atom, this does not yield a stable structure.

The general formula for diketopiperazine and its analogs is shown below.

25



30

Wherein n is between 0 and 7, Q is, independently, a C₁₋₂₀ straight,

branched or cyclic alkyl, aralkyl, alkaryl, alkenyl, alkynyl, heteroalkyl, heterocyclic, alkyl-heterocyclic, or heterocyclic-alkyl; T is -C(O)O, -OC(O), -C(O)NH, -NH, -NQ, -OQO, -O, -NHC(O), -OP(O), -P(O)O, -OP(O)₂, -P(O)₂O, -OS(O)₂, or -S(O)₃; U is an acid group, such as a

5 carboxylic acid, phosphoric acid, phosphonic acid and sulfonic acid, or a basic group, such as primary, secondary and tertiary amines, quaternary ammonium salts, guanidine, aniline, heterocyclic derivatives, such as pyridine and morpholine, or a zwitterionic C₁₋₂₀ chain containing at least one acidic group and at least one basic group, for example, those described

10 above, wherein the side chains can be further functionalized with an alkene or alkyne group at any position, one or more of the carbons on the side chain can be replaced with an oxygen, for example, to provide short polyethylene glycol chains, one or more of the carbons can be functionalized with an acidic or basic group, as described above, and wherein the ring atoms X at

15 positions 1 and 4 are either O or N.

As used herein, "side chains" are defined as Q-T-Q-U or Q-U, wherein Q, T, and U are defined above.

Examples of acidic side chains include, but are not limited, to cis and trans -CH=CH-CO₂H, -CH(CH₃)=CH(CH₃)-CO₂H,

20 -(CH₂)₃-CO₂H, -CH₂CH(CH₃)-CO₂H, -CH(CH₂CO₂H)=CH₂, -(tetrafluoro)benzoic acid, -benzoic acid and -CH(NHC(O)CF₃)-CH₂-CO₂H.

Examples of basic side chains include, but are not limited to, -aniline, -phenyl-C(NH)NH₂, -phenyl-C(NH)NH(alkyl),

25 -phenyl-C(NH)N(alkyl)₂ and -(CH₂)₄NHC(O)CH(NH₂)CH(NH₂)CO₂H.

Examples of zwitterionic side chains include, but are not limited to, -CH(NH₂)-CH₂-CO₂H and -NH(CH₂)₁₋₂₀CO₂H.

The term aralkyl refers to an aryl group with an alkyl substituent.

The term heterocyclic-alkyl refers to a heterocyclic group with an

30 alkyl substituent.

The term alkaryl refers to an alkyl group that has an aryl substituent.

The term alkyl-heterocyclic refers to an alkyl group that has a heterocyclic substituent.

The term alkene, as referred to herein, and unless otherwise specified, refers to an alkene group of C₂ to C₁₀, and specifically includes vinyl and allyl.

The term alkyne, as referred to herein, and unless otherwise specified, refers to an alkyne group of C₂ to C₁₀.

As used herein, "diketopiperazines" includes diketopiperazines and derivatives and modifications thereof falling within the scope of the above-general formula.

Fumaryl diketopiperazine is most preferred for pulmonary applications.

(ii). Synthesis

Diketopiperazines can be formed by cyclodimerization of amino acid ester derivatives, as described by Katchalski, *et al.*, *J. Amer. Chem. Soc.* 68:879-80 (1946), by cyclization of dipeptide ester derivatives, or by thermal dehydration of amino acid derivatives in high-boiling solvents, as described by Kopple, *et al.*, *J. Org. Chem.* 33(2):862-64 (1968). 2,5-diketo-3,6-di(aminobutyl)piperazine (Katchalski *et al.* refer to this as lysine anhydride) was prepared via cyclodimerization of N-epsilon-P-L-lysine in molten phenol, similar to the Kopple method in *J. Org. Chem.*, followed by removal of the blocking (P)-groups with 4.3 M HBr in acetic acid. This route is preferred because it uses a commercially available starting material, it involves reaction conditions that are reported to preserve stereochemistry of the starting materials in the product and all steps can be easily scaled up for manufacture.

Diketomorpholine and diketooxetane derivatives can be prepared by stepwise cyclization in a manner similar to that disclosed in Katchalski, *et al.*, *J. Amer. Chem. Soc.* 68:879-80 (1946).

Diketopiperazines can be radiolabelled. Means for attaching radiolabels are known to those skilled in the art. Radiolabelled diketopiperazines can be prepared, for example, by reacting tritium gas with

those compounds listed above that contain a double or triple bond. A carbon-14 radiolabelled carbon can be incorporated into the side chain by using ^{14}C labeled precursors which are readily available. These radiolabelled diketopiperazines can be detected *in vivo* after the resulting microparticles are administered to a subject.

(a) Synthesis of Symmetrical

Diketopiperazine Derivatives

The diketopiperazine derivatives are symmetrical when both side chains are identical. The side chains can contain acidic groups, basic groups, or combinations thereof.

One example of a symmetrical diketopiperazine derivative is 2,5-diketo-3,6-di(4-succinylaminobutyl)piperazine. 2,5-diketo-3,6-di(aminobutyl) piperazine is exhaustively succinylated with succinic anhydride in mildly alkaline aqueous solution to yield a product which is readily soluble in weakly alkaline aqueous solution, but which is quite insoluble in acidic aqueous solutions. When concentrated solutions of the compound in weakly alkaline media are rapidly acidified under appropriate conditions, the material separates from the solution as microparticles.

Other preferred compounds can be obtained by replacing the succinyl group(s) in the above compound with glutaryl, maleyl or fumaryl groups.

(b) Synthesis of Asymmetrical

Diketopiperazine Derivatives

One method for preparing unsymmetrical diketopiperazine derivatives is to protect functional groups on the side chain, selectively deprotect one of the side chains, react the deprotected functional group to form a first side chain, deprotect the second functional group, and react the deprotected functional group to form a second side chain.

Diketopiperazine derivatives with protected acidic side chains, such as cyclo-Lys(P)Lys(P), wherein P is a benzyloxycarbonyl group, or other protecting group known to those skilled in the art, can be selectively deprotected. The protecting groups can be selectively cleaved by using limiting reagents, such as HBr in the case of the benzyloxycarbonyl group, or fluoride ion in the case of silicon protecting groups, and by using controlled

time intervals. In this manner, reaction mixtures which contain unprotected, monoprotected and di-protected diketopiperazine derivatives can be obtained. These compounds have different solubilities in various solvents and pH ranges, and can be separated by selective precipitation and removal.

- 5 An appropriate solvent, for example, ether, can then be added to such reaction mixtures to precipitate all of these materials together. This can stop the deprotection reaction before completion by removing the diketopiperazines from the reactants used to deprotect the protecting groups. By stirring the mixed precipitate with water, both the partially and
- 10 completely reacted species can be dissolved as salts in the aqueous medium. The unreacted starting material can be removed by centrifugation or filtration. By adjusting the pH of the aqueous solution to a weakly alkaline condition, the asymmetric monoprotected product containing a single protecting group precipitates from the solution, leaving the completely
- 15 deprotected material in solution.

- In the case of diketopiperazine derivatives with basic side chains, the basic groups can also be selectively deprotected. As described above, the deprotection step can be stopped before completion, for example, by adding a suitable solvent to the reaction. By carefully adjusting the solution pH, the
- 20 deprotected derivative can be removed by filtration, leaving the partially and totally deprotected derivatives in solution. By adjusting the pH of the solution to a slightly acidic condition, the monoprotected derivative precipitates out of solution and can be isolated.

- Zwitterionic diketopiperazine derivatives can also be selectively
- 25 deprotected, as described above. In the last step, adjusting the pH to a slightly acidic condition precipitates the monoprotected compound with a free acidic group. Adjusting the pH to a slightly basic condition precipitates the monoprotected compound with a free basic group.

- Limited removal of protecting groups by other mechanisms,
- 30 including but not limited to cleaving protecting groups that are cleaved by hydrogenation by using a limited amount of hydrogen gas in the presence of palladium catalysts. The resulting product is also an asymmetric partially

deprotected diketopiperazine derivative. These derivatives can be isolated essentially as described above.

The monoprotected diketopiperazine is reacted to produce a diketopiperazine with one sidechain and protecting group. Removal of
5 protecting groups and coupling with other side chains yields unsymmetrically substituted diketopiperazines with a mix of acidic, basic, and zwitterionic sidechains.

Other materials that exhibit this response to pH can be obtained by functionalizing the amide ring nitrogens of the diketopiperazine ring.

10 C. Transport Enhancers

In a preferred embodiment, the active agent is complexed with a transport enhancer which is degradable and capable of forming hydrogen bonds with the target biological membrane in order to facilitate transport of the agent across the membrane. The transport enhancer also is capable of
15 forming hydrogen bonds with the active agent, if charged, in order to mask the charge and facilitate transport of the agent across the membrane. A preferred transport enhancer is diketopiperazine.

The transport enhancer preferably is biodegradable and may provide linear, pulsed or bulk release of the active agent. The transport enhancer
20 may be a natural or synthetic polymer and may be modified through substitutions or additions of chemical groups, including alkyl, alkylene, hydroxylations, oxidations, and other modifications routinely made by those skilled in the art.

A preferred transport enhancer is fumaryl diketopiperazine. Other
25 diketopiperazines which may be useful as a transport enhancer are described above.

Like most proteins and peptides, insulin is a charged molecule, which impedes its ability to cross charged biological membranes. It has been found that when insulin hydrogen bonds to fumaryl diketopiperazine, the charge of
30 the peptide is masked, thereby facilitating or enhancing the passage of insulin across the membranes, such as mucosal membranes, and into the blood.

II. Methods

A. Encapsulation

In one embodiment, active agent is encapsulated within microparticles by dissolving a diketopiperazine with acidic side chains in bicarbonate or other basic solution, adding the active agent in solution or suspension, and then precipitating the microparticle by adding acid, such as 1 M citric acid.

In another embodiment, active agent is encapsulated within microparticles by dissolving a diketopiperazine with basic side chains in an acidic solution, such as 1 M citric acid, adding the active agent in solution or suspension, and then precipitating the microparticle by adding bicarbonate or another basic solution.

In still another embodiment, active agent is encapsulated within microparticles by dissolving a diketopiperazine with both acidic and basic side chains in an acidic or basic solution, adding the active agent in solution or suspension to be encapsulated, then precipitating the microparticle by neutralizing the solution.

The microparticles can be stored in the dried state and suspended for administration to a patient. In the first embodiment, the reconstituted microparticles maintain their stability in an acidic medium and dissociate as the medium approaches physiological pH in the range of between 6 and 14. In the second embodiment, suspended microparticles maintain their stability in a basic medium and dissociate at a pH of between 0 and 6. In the third embodiment, the reconstituted microparticles maintain their stability in an acidic or basic medium and dissociate as the medium approaches physiological pH in the range of pH between 6 and 8.

The impurities typically are removed when the microparticles are precipitated. However, impurities also can be removed by washing the particles to dissolve the impurities. A preferred wash solution is water or an aqueous buffer. Solvents other than water also can be used to wash the microspheres or precipitate the diketopiperazines, in order to remove impurities that are not water soluble. Any solvent in which neither the cargo

nor the fumaryl diketopiperazine is soluble are suitable. Examples include acetic acid, ethanol, and toluene.

In an alternative embodiment, microparticles of diketopiperazine are prepared and provided in a suspension, typically an aqueous suspension, to which a solution of the active agent then is added. The suspension is then lyophilized or freeze dried to yield diketopiperazine microparticles having a coating of active agent. In a preferred embodiment, the active agent is insulin in a hexameric form. Zinc ions can then be removed by washing the microparticles with an appropriate solvent.

As used herein, the term "entrapped" with reference to an active agent in/with a diketopiperazine includes coating of the active agent onto microparticles of the diketopiperazine.

The diketopiperazine microparticles have been found to have a higher affinity for insulin than does zinc. Insulin has been found to be stabilized within an ordered lattice array of fumaryl diketopiperazine. In this state, in the sufficient absence of zinc ions, the insulin is predominately dimeric and monomeric, as opposed to its hexameric state. The insulin therefore more readily dissociates to its monomeric state, which is the state in which insulin exerts its biological activity.

Other complexing agents may be substituted for the diketopiperazine. Other representative complexing agents include serum albumin and other proteins, alginic acid, antibodies, cyclodextrins, phospholipids, and lecithin. For example, insulin contaminated with zinc can be complexed with bovine serum albumin. The complex can be dialyzed in tubing with a molecular weight cut-off below 1,000 Daltons to separate and remove the zinc. Once sufficient amounts of zinc have been dialyzed away, as evidenced by its presence in the dialysate, the dispersion is transferred to dialysis tubing with a molecular weight cut-off below 10,000 Daltons. Only monomeric insulin will pass through the tubing into the dialysate, leaving any remaining hexameric zinc complexed insulin behind. The purified insulin can be captured from the dialysate.

These materials may not, however, provide sufficient stabilization of unstable or labile drugs.

B. Administration

The compositions of active agent described herein can be
5 administered to patients in need of the active agent. The compositions preferably are administered in the form of microparticles, which can be in a dry powder form for pulmonary administration or suspended in an appropriate pharmaceutical carrier, such as saline.

The microparticles preferably are stored in dry or lyophilized form
10 until immediately before administration. The microparticles then can be administered directly as a dry powder, such as by inhalation using, for example, dry powder inhalers known in the art. Alternatively, the microparticles can be suspended in a sufficient volume of pharmaceutical carrier, for example, as an aqueous solution for administration as an aerosol.

15 The microparticles also can be administered via oral, subcutaneous, and intravenous routes.

The compositions can be administered to any targeted biological membrane, preferably a mucosal membrane of a patient. In a preferred embodiment, the patient is a human suffering from Type II diabetes. In a
20 preferred embodiment, the composition delivers insulin in biologically active form to the patient, which provides a spike of serum insulin concentration which simulates the normal response to eating.

In a preferred embodiment, hexameric insulin is entrapped in fumaryl diketopiperazine to form a solid precipitate of monomeric insulin in the
25 fumaryl diketopiperazine, which then is washed with aqueous solution to remove the free zinc. This formulation demonstrates blood uptake following pulmonary administration at a rate 2.5 times the rate of insulin uptake following subcutaneous injection, with peak blood levels occurring at between 7.5 and 10 minutes after administration.

30 The range of loading of the drug to be delivered is typically between about 0.01% and 90%, depending on the form and size of the drug to be delivered and the target tissue. In a preferred embodiment using

diketopiperazines, the preferred range is from 0.1% to 50% loading by weight of drug. The appropriate dosage can be determined, for example, by the amount of incorporated/encapsulated agent, the rate of its release from the microparticles, and, in a preferred embodiment, the patient's blood
5 glucose level.

One preferred application is in the treatment of hyperinsulinemia. In a preferred embodiment, microparticles of the composition wherein the active agent is glucagon can be administered by continuous subcutaneous infusion. Glucagon is an extremely unstable peptide, but can be stabilized in
10 particles of diketopiperazine, for example. The stabilized glucagon/diketopiperazine microparticles can be made by adding glucagon to a solution of diketopiperazine which hydrogen bonds to the glucagon and when the solution is acidified, such as by adding a food acid, both the diketopiperazine and the glucagon self-assemble to form uniform
15 microspheres having a mean particle size of, for example, about 2 μm . In this process, approximately 95% of the glucagon is pulled out of solution and is evenly distributed within the diketopiperazine microparticle. These particles can readily be suspended and infused subcutaneously with a standard infusion pump. Then the glucagon/diketopiperazine particles are
20 contacted with the near neutral pH environment of the subcutaneous fluid, where they dissolve, thereby releasing glucagon in its pharmacologically active state.

The compositions and methods described herein are further described by the following non-limiting examples.

25 **Example 1: Removal of Zinc From U.S.P. Injectable Insulin**

Insulin trapped in fumaryl diketopiperazine was analyzed to assess whether zinc was removed during the entrapment process. The insulin used as the starting material met U.S.P. standards for injectable insulin, and according to the certificate of analysis, the insulin contained a considerable
30 quantity of zinc: 0.41%. This insulin was then entrapped in fumaryl diketopiperazine to form a solid fumaryl diketopiperazine/insulin mixture, as described above.

Following entrapment of the insulin in fumaryl diketopiperazine, the amount of zinc theoretically should be present in the same proportion as it existed in the neat insulin. Using the certificate of analysis value, it was calculated that one should expect to find 697 parts per million (ppm) of zinc per gram in the solid yield of fumaryl diketopiperazine/insulin. Surprisingly, the quantity of zinc present the solid fumaryl diketopiperazine/insulin was measured to be only 6 ppm. The “missing” zinc was presumably eliminated with the water used to wash the insulin/fumaryl diketopiperazine precipitate.

Example 2: Bioavailability of Insulin in Diketopiperazine

Pulmonary Formulation

Subjects and Methods

The study was reviewed and approved by the ethical review committee of the Heinrich-Heine-University, Düsseldorf, and conducted according to local regulations, the Declaration of Helsinki and the rules of Good Clinical Practice.

The study was conducted with 5 healthy male volunteers. Inclusion criteria were good health, as judged by physical examination, age: 18 to 40 years, body mass index: 18 to 26 kg/m², capability to reach peak inspiratory flow of 4 l/sec measured by a computer assisted spirometry and a FEV₁ equal to or greater than 80% of predicted normal (FEV₁ = forced expiratory volume in one second). Exclusion criteria were Diabetes mellitus type 1 or 2, prevalence of human insulin antibodies, history of hypersensitivity to the study medication or to drugs with similar chemical structures, history or severe or multiple allergies, treatment with any other investigational drug in the last 3 months before study entry, progressive fatal disease, history of drug or alcohol abuse, current drug therapy with other drugs, history significant cardiovascular, respiratory, gastrointestinal, hepatic, renal, neurological, psychiatric and/or hematological disease, ongoing respiratory tract infection or subjects defined as being smokers with evidence or history of tobacco or nicotine use.

Conduct of the Study

On the morning of the study days, the subjects came to the hospital

(fasting, except for water, from midnight onward) at 7:30 a.m. The subjects were restricted from excessive physical activities and an intake of alcohol for 24 hours before each treatment day. They were randomly assigned to one of the three treatment arms. The subjects received a constant intravenous regular human insulin infusion, which was kept at $0.15 \text{ mU min}^{-1} \text{ kg}^{-1}$ so that serum insulin concentrations were established at 10-15 U/ml during a period of 2 hours before time point 0. This low-dose infusion was continued throughout the test to suppress endogenous insulin secretion. Blood glucose was kept constant at a level of 90 mg/dl throughout the glucose clamp by a glucose controlled infusion system (BIOSTATORTM). The glucose clamp algorithm was based on the actual measured blood glucose concentration and the grade of variability in the minutes before to calculate the glucose infusion rates for keeping the blood glucose concentration constant. The insulin application (5 U i.v. or 10 U s.c. injection or three deep breaths inhalation per capsule (2 capsules with 50 U each) applied with a commercial inhalation device (Boehringer Ingelheim)) had to be finished immediately before time point 0. The duration of the clamp experiment was 6 hours from time point 0. Glucose infusion rates, blood glucose, serum-insulin and C-peptide were measured.

20 Bioefficacy and Bioavailability

To determine bioefficacy, the areas under the curve of the glucose infusion rates were calculated for the first 3 hours (AUC_{0-180}) after the administration and for the overall observation period of six hours after the administration (AUC_{0-360}) and were correlated to the amount of insulin applied. To determine bioavailability, the areas under the curve of the insulin concentrations were calculated for the first 3 hours (AUC_{0-180}) after the administration and for the overall observation period of six hours after the administration (AUC_{0-360}) and correlated to the amount of insulin applied.

In this clamp study, inhalation of 100 U of TECHNOSPHERETM/Insulin was well tolerated and was demonstrated to have a substantial blood glucose lowering effect with a relative bioavailability of 25.8% for the first three hours as calculated from the

achieved serum insulin concentrations. TECHNOSPHERES™ are microparticles (also referred to herein as microspheres) formed of diketopiperazine that self-assembles into an ordered lattice array at particular pHs, typically a low pH. They typically are produced to have a mean diameter between about 1 and about 5 μm .

Results

The pharmacokinetic results are illustrated in Figures 1 and 2 and in Table 1.

Efficacy Results

Inhalation of 100 U of TECHNOSPHERE™/Insulin (inhalation of 100 U) revealed a peak of insulin concentration after 13 min (intravenous (i.v.) (5U): 5 min, subcutaneous (s.c.) (10 U): 121 min) and a return of the insulin levels to baseline after 180 min (i.v.: 60 min, s.c. 360 min). Biological action as measured by glucose infusion rate peaked after 39 min (i.v. 14 min, s.c.: 163 min) and lasted for more than 360 min (i.v.: 240 min, s.c.: > 360 min). Absolute bioavailability (comparison to i.v. application) was $14.6 \pm 5.1\%$ for the first 3 hours and $15.5 \pm 5.6\%$ for the first 6 hours. Relative bioavailability (comparison to s.c. application) was $25.8 \pm 11.7\%$ for the first 3 hours and $16.4 \pm 7.9\%$ for the first 6 hours.

Table 1: Pharmacokinetic Parameters

<u>Parameter Calculated on Glucose Infusion Rate</u>	<u>Intravenous Administration</u>	<u>Inhaled</u>	<u>Subcutaneous Administration</u>
T50%*	9 min	13 min	60 min
Tmax	14 min	39 min	163 min
T-50%**	82 min	240 min	240 min
T to baseline	240 min	>360 min	>360 min
<u>Parameter Calculated on Insulin Levels</u>	<u>Intravenous Administration</u>	<u>Inhaled</u>	<u>Subcutaneous Administration</u>
T50%*	2 min	2.5 min	27 min
Tmax	5 min	13 min	121 min
T-50%**	6 min	35 min	250 min
T to baseline	60 min	180 min	360 min

* time from baseline to half-maximal values

** time from baseline to half-maximal after passing Tmax

Safety Results

TECHNOSPHERE™/Insulin was shown to be safe in all patients.

One patient was coughing during the inhalation without any further symptoms or signs of deterioration of the breathing system.

Conclusions

Inhalation of 100 U of TECHNOSPHERE™/Insulin was well tolerated and was demonstrated to have a substantial blood glucose lowering effect with a relative bioavailability of 25.8% for the first 3 hours as calculated from the achieved serum insulin concentrations.

Summary

In this study, the inhalation of TECHNOSPHERE™/Insulin (the formulation of example 1) was demonstrated in healthy human subjects to have a time-action profile with a rapid peak of insulin concentration (Tmax: 13 min) and rapid onset of action (Tmax: 39 min) and a sustained action over more than 6 hours. The total metabolic effect measured after inhalation of 100 U of TECHNOSPHERE™/Insulin was larger than after subcutaneous injection of 10 U of insulin. The relative bioefficacy of TECHNOSPHERE™/Insulin was calculated to be 19.0%, while the relative bioavailability was determined to be 25.8% in the first three hours.

The data also show that inhalation of TECHNOSPHERE™/Insulin resulted in a much more rapid onset of action than s.c. insulin injection that was close to the onset of action of i.v. insulin injection, while duration of action of TECHNOSPHERE™/Insulin was comparable to that of s.c. insulin injection.

The drug was well tolerated and no serious adverse events were reported during the entire trial.

Example 3: Removal of Impurity From Proprietary Peptide

A proprietary peptide containing an impurity was trapped in fumaryl diketopiperazine, forming a peptide/fumaryl diketopiperazine precipitate. The precipitate was washed with water to remove the impurity. The peptide is rather unstable and trapping it in fumaryl diketopiperazine markedly improves its stability; both as a dry powder and in aqueous suspension for injection.

Example 4: Stabilized Glucagon Formulations

Formulation

Glucagon was formulated under sterile conditions, into a stabilized complex by precipitation in acidic solution with fumaryl diketopiperazine (3,6-bis[N-fumaryl-N-(n-butyl)amino]-2,5-diketopiperazine). The complex was washed and lyophilized, yielding a sterile dry powder formulation of diketopiperazine/glucagon (hereinafter referred to as "TG") containing from 1.2 to 8.2% glucagon by weight, depending upon the formulation parameters desired (allowing physicians to increase dose yet keep the volume constant). The TG powder was suspended in an appropriate media suitable for subcutaneous delivery in a MiniMed 507C infusion pump.

Stability Protocol

Glucagon and TG were suspended in infusion media and incubated at 40 °C in a water bath for varying amounts of time up to 150 hours.

Glucagon HPLC Analysis

An adaptation of USP method for glucagon analysis was employed. A Waters Symmetry Shield RP8 column (5 µm, 3.9 x 150 mm) and guard RP8 column (5 µm, 3.9 x 20 mm) were used at a flow rate of 1 mL/min. and

a detection wavelength of 214 nm. The gradient method consisted of mobile phase A: 9.8 g NaH₂PO₄ (0.0816 M) and 170 mg L-cysteine (1.4 mM) per liter HPLC grade water, adjusted pH to 2.6 with phosphoric acid; and B: acetonitrile. Glucagon solutions were diluted as needed with water and injected. TG samples were prepared by adding 1/10th volume 1 M Tris pH 10.0 to sample to solubilize the fumaryl diketopiperazine.

Rat Study Protocol

Sprague Dawley rats 200-250 g were fasted overnight and given subcutaneous injection of glucagon or TG (0.75 mg/kg) in an appropriate media that had been held at 25 °C for 0, 24, or 48 hours. Blood samples were taken at -10, -5, 0, 5, 10, 15, 20, 30, 45, and 60 minutes post dose and analyzed for blood glucose (HemCue B-glucose analyzer, Hemocue AB, Angelholm Sweden). Mean baseline was determined (pre-dose measurements) and was subtracted from the subsequent data and plotted vs. time. This was done to assure that the TG formulation, which appeared to not degrade significantly, showed appropriate pharmacological activity.

Results

Following 40 °C incubation, HPLC analysis showed an increase in breakdown products in the glucagon preparation. By contrast, TG has only one minor degradation peak (RT=6) which correlated with the slightly less active oxidative form of glucagon. Glucagon without diketopiperazine (i.e. without TECHNOSPHERESTM) had many degradation peaks, some of which contributed to an enhanced effect and others that reduced the potency of glucagon.

The TG sterile lyophilized powder was shipped frozen to a hospital, where it was re-suspended in sterile media. The material re-suspended well and each vial was continuously infused over a 72 hour period.

Conclusion

Standard preparations of glucagon are not suitable for regulation of blood glucose by continuous subcutaneous infusion. Administration of such preparations containing variable amounts of the deamidated and hydrolysed forms resulted in highly variable blood glucose levels. Suspensions of

TECHNOSPHERESTM/glucagon, which is stabilized, does not aggregate and contains clinically irrelevant amounts of breakdown products. As such TG can be and has been used as a therapy for hyperinsulinemia, providing consistent, elevated glucose levels when administered subcutaneously over

5 time.



We claim:

1. A method for purifying an active agent comprising
providing an active agent containing an impurity to be removed;
entrapping the active agent in a diketopiperazine to form a mixture;
and
removing essentially all of the impurity from the mixture.
2. The method of claim 1 wherein the active agent is a peptide or protein.
3. The method of claim 2 wherein the peptide or protein is selected from the group consisting of insulin, salmon calcitonin, parathyroid hormone 1-34, octreotide, leuprolide, and RSV peptide.
4. The method of claim 3 wherein the active agent is insulin.
5. The method of claim 1 wherein the impurity is a multi-valent ion.
6. The method of claim 4 wherein the active agent of step (a) is an insulin complex and the impurity is a zinc ion.
7. The method of claim 6 wherein the complex is hexameric insulin.
8. The method of any of claims 1-7 wherein the diketopiperazine is fumaryl diketopiperazine.
9. The method of any of claims 1-8 wherein the active agent is entrapped by a process which comprises
making a solution of the diketopiperazine,
making a solution or suspension of the active agent,
combining the diketopiperazine solution with the solution or suspension of active agent, and
precipitating microparticles of the diketopiperazine in which the active agent is dispersed, thereby forming the mixture.
10. The method of claim 9 further comprising washing the microparticles with an aqueous solvent for the impurity.
11. A composition for the administration of a peptide to a patient, comprising a peptide stabilized in a diketopiperazine.

12. The composition of claim 11 wherein the peptide is dimeric or monomeric insulin.
13. The composition of claim 11 wherein the peptide is glucagon.
14. The composition of any of claims 11-13 wherein the diketopiperazine is fumaryl diketopiperazine.
15. The composition of claim 12 substantially free of zinc ions.
16. The composition of any of claims 11-15 wherein the peptide is dispersed within or coated onto microparticles of the diketopiperazine.
17. The composition of claim 16 wherein the microparticles are provided in the form of a dry powder.
18. The composition of claim 16 wherein the microparticles are provided as an aqueous suspension in a pharmaceutically acceptable carrier.
19. The composition of any of claims 11, 15, or 17 which is in a form suitable for pulmonary administration.
20. The composition of claim 11 made by the methods of any of claims 1-10.
21. A method for administering an active agent to a mucosal membrane of a person in need of the active agent, comprising
administering to the mucosal membrane a composition which comprises
microparticles formed of (i) the active agent and (ii) an effective amount of a transport enhancer to facilitate transport of the active agent across the mucosal membrane by masking a charge of the agent, if any, by hydrogen bonding the transport enhancer to the mucosal membrane, or a combination thereof.
22. The method claim 21 wherein the active agent is a charged molecule.
23. The method of claim 22 wherein the active agent is insulin.
24. The method claim 21 wherein the transport enhancer forms hydrogen bonds with the active agent to mask its charge.
25. The method of claim 21 wherein the transport enhancer is fumaryl diketopiperazine.


26. The method of any of claims 21-25 wherein the composition is administered to the lungs via inhalation.
27. A method for delivering insulin to a patient in need thereof, comprising
administering to the patient an effective amount of the insulin in a composition which comprises microparticles of a diketopiperazine in which monomeric insulin is encapsulated.
28. The method of claim 27 wherein the diketopiperazine is fumaryl diketopiperazine.
29. The method of claim 28 wherein the composition is in a dry powder form administered to the lungs via inhalation.
30. The method of any of claims 27-29 wherein the patient is a Type II diabetic.
31. The method of claim 30 wherein the composition is administered concurrently with, or less than about 20 minutes prior to, the patient eating a meal.
32. The method of claim any of claims 27-31 wherein the composition is provided in one or more unit doses of insulin, each dose equivalent to about 6 IU of insulin.

1/2


**FIG. 1A****FIG. 1B**

2/2

glucose infusion rate (mg/kg/min)

**FIG. 2A**

INSULIN CONCENTRATIONS

**FIG. 2B**

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/17984

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K1/36 C07D241/08 //A61K47/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 503 852 A (STEINER SOLOMON S ET AL) 2 April 1996 (1996-04-02)	1-7, 9-13, 15-24, 26, 27, 29-32
Y	claim 13, claims 8/9	8, 14, 25, 28
Y	WO 96 36314 A (PHARMACEUTICAL DISCOVERY CORP) 21 November 1996 (1996-11-21) claims	1-28
A	EP 0 220 958 A (LILLY CO ELI) 6 May 1987 (1987-05-06) whole document	1-28
	--- -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

19 December 2000

Date of mailing of the international search report

08/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Kronester-Frei, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/17984

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GUPTA P K ET AL: "CONTEMPORARY APPROACHES IN AEROSOLIZED DRUG DELIVERY TO THE LUNG" JOURNAL OF CONTROLLED RELEASE, NL, ELSEVIER SCIENCE PUBLISHERS B.V. AMSTERDAM, vol. 17, no. 2, 1 October 1991 (1991-10-01), pages 127-147, XP000232469 ISSN: 0168-3659 whole document	1-28
P, X	--- DATABASE CAPLUS 'Online! LIAN H ET AL: "A self-complementary, self-assembling microsphere system: application for intravenous delivery of the antiepileptic and neuroprotectant compound felbamate." retrieved from STN Database accession no. 2000:522278 XP002154327	1-7, 9-13, 15-24, 26, 27, 29-32
P, Y	abstract abstract & J. PHARM. SCI., (2000). VOL. 89, NO. 7, PP. 867-875. CODEN: JPMSAE. ISSN. 0022-3549., Pharmaceutical Discovery Corporation, Elmsford	8, 14, 25, 28
P, Y	--- WO 99 52506 A (BENGIS HOLGER ;GRANDE JUERGEN (DE); AVENTIS RES & TECH GMBH & CO (D) 21 October 1999 (1999-10-21) whole document -----	1-28

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/17984

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5503852 A	02-04-1996	US 5352461 A AT 132744 T AU 680408 B AU 3804493 A CA 2131366 A DE 69301311 D DE 69301311 T DK 630236 T EP 0630236 A ES 2089844 T GR 3019622 T JP 2617273 B JP 7506818 T WO 9318754 A	04-10-1994 15-01-1996 31-07-1997 21-10-1993 30-09-1993 22-02-1996 22-08-1996 03-06-1996 28-12-1994 01-10-1996 31-07-1996 04-06-1997 27-07-1995 30-09-1993
WO 9636314 A	21-11-1996	AU 5740996 A US 6071497 A	29-11-1996 06-06-2000
EP 0220958 A	06-05-1987	AU 6441486 A CN 86107569 A DK 512286 A HU 41811 A JP 62106100 A KR 8904352 B NZ 218088 A PT 83613 A, B SU 1531858 A US 4782139 A ZA 8608168 A	30-04-1987 08-07-1987 29-04-1987 28-05-1987 16-05-1987 31-10-1989 27-01-1989 01-11-1986 23-12-1989 01-11-1988 29-06-1988
WO 9952506 A	21-10-1999	AU 3421099 A	01-11-1999